



Declawing Facts

KNOW THE TRUTH



Declawing = amputation of cat's toe at first joint

(like cutting off a human's first knuckle)

33%

of cats suffer at least one behavioral problem after declaw surgery



Illegal in

37

Countries

+ 8 U.S. Cities

Declawing can produce:
Chronic pain
Arthritis/lameness
Nerve/tissue damage
Gangrene
Abscesses
Paw pad atrophy
Bone regrowth

18%
show increased biting

17%
suffer wounds re-opening

15%
will not use the litterbox

11%
suffer from lameness

10%
see nail regrowth

up to **50%**

of declawed cats develop acute physical complications



up to **20%**

of declawed cats develop long-term physical complications

"The pain must be excruciating; it is certainly a **gruesome spectacle** to watch."
--A Vet Tech



75-90% of pet owners change their mind about declawing once they know the risks

"Claws are a physically, socially, and emotionally vital part of every cat."

--Jackson Galaxy, Cat Behaviorist



"Unlike routine recoveries... declawing surgery results in cats bouncing off the walls of the recovery cage because of **excruciating pain...**"

--Dr. Nicholas Dodman, DVM

Litterbox issues are

2x

more common in declawed cats

Osteomyelitis affects

30%

of declawed cats (a painful bone infection)

You wouldn't declaw a dog... you trim his nails



Sources:

JAVMA (Journal of the American Veterinary Medical Association), AVMA.org, Veterinary Medicine, DVM Best Practices, James Gaynor, DVM, Colorado State University Veterinary Medical School, pawproject.org, pet360.com, littlebigcat.com, Vet Forum, paw-rescue.org, moggies.co.uk, declaw.lisaviolet.com, maxshouse.com, World Small Animal Veterinary Association, jacksongalaxy.com